

ACL of 150 mt is based on a rebuilding plan with a target year to rebuild of 2051 and an SPR harvest rate of 86.4 percent. 16.5 mt is deducted from the ACL for the Tribal fishery (10.9 mt), open access fishery (0.4 mt) and research catch (5.2 mt), resulting in a fishery HG of 133.5 mt.

ddd/ Pacific whiting. Pacific whiting are assessed annually. The final specifications will be determined consistent with the U.S.-Canada Pacific Whiting Agreement and will be announced after the Council's April 2013 meeting.

eee/ Petrale sole. A petrale sole stock assessment was prepared for 2011. In 2011 the petrale sole stock was estimated to be at 18 percent of its unfished biomass. The OFL of 2,711 mt is based on the 2011 assessment with an $F_{30\%} F_{MSY}$ proxy. The ABC of 2,592 mt is a 4 percent reduction from the OFL ($\sigma=0.36/P^*=0.45$) as it's a category 1 stock. The ACL is set equal to the ABC. 234 mt is deducted from the ACL for the Tribal fishery (220 mt), the incidental open access fishery (2.4 mt), and research catch (11.6 mt), resulting in a fishery HG of 2,358 mt.

fff/ Sablefish north. A coastwide sablefish stock assessment was prepared in 2011. The coastwide sablefish biomass was estimated to be at 33 percent of its unfished biomass in 2011. The coastwide OFL of 6,621 mt is based on the 2011 stock assessment with an F_{MSY} proxy of $F_{45\%}$. The coastwide ABC of 6,045 mt is an 8.7 percent reduction from the OFL ($\sigma=0.36/P^*=0.40$). The 40-10 harvest policy was applied to the ABC to derive a coastwide ACL value. Then the ACL value was apportioned, north and south of 36° N. lat., using the average of annual swept area biomass (2003-2010) from the NMFS NWFSC trawl survey, between the northern and southern areas with 73.6 percent going to the area north of 36° N. lat. and 26.4 percent going to the area south of 36° N. lat. The northern ACL is 4,012 mt and is reduced by 401 mt for the tribal allocation (10 percent of the ACL north of 36° N. lat.). The 401 mt Tribal allocation is