gg/ Sablefish south. The ACL for the area south of 36° N. lat. is 1,560 mt (26.4 percent of the calculated coastwide ACL value). 5 mt is deducted from the ACL for the incidental open access fishery (2 mt) and research catch (3 mt), resulting in a fishery HG of 1,555 mt.

hh/ Shortbelly rockfish. A non quantitative assessment was conducted in 2007. The spawning stock biomass of shortbelly rockfish was estimated at 67 percent of its unfished biomass in 2005. The OFL of 6,950 mt was recommended for the stock in 2014 with an ABC of 5,789 mt (σ=0.72 with a P* of 0.40). The 50 mt ACL is slightly higher than recent landings and is in recognition of the stock's importance as a forage species in the California Current ecosystem. 2 mt is deducted from the ACL for research catch, resulting in a fishery HG of 48 mt.

ii/ Shortspine thornyhead. A coastwide stock assessment was conducted in 2005 and the stock was estimated to be at 63 percent of its unfished biomass in 2005. A coastwide OFL of 2,310 mt is based on the 2005 stock assessment with a $F_{50\%}$ F_{MSY} proxy. The coastwide ABC of 2,208 mt is a 4 percent reduction from the OFL (σ=0.36/P*=0.45) as it's a category 1 stock. For the portion of the stock that is north of 34°27' N. lat., the ACL is 1,525 mt. The northern ACL is 66 percent of the coastwide OFL for the portion of the biomass found north of 34°27' N. lat. 59.22 mt is deducted from the ACL for the Tribal fishery (50 mt), the incidental open access fishery (2 mt), and research catch (7.22mt) resulting in a fishery HG of 1,466 mt for the area north of 34°27' N. lat. For that portion of the stock south of 34°27' N. lat. the ACL is 393 mt which is 34 percent of the coastwide OFL for the portion of the biomass found south of 34°27' N. lat. reduced by 50 percent as a precautionary adjustment. 42 mt is deducted from the ACL for the incidental open access fishery (41 mt), and research catch (1 mt) resulting in a fishery HG of 351 mt for the area south of 34°27' N. lat.