

Table 1b. to Part 660, Subpart C – 2015, Allocations by Species or Species Group. (Weights in Metric Tons)

Species	Area	Fishery HG or ACT	Trawl		Non-trawl	
			%	Mt	%	Mt
BOCACCIO a/	S of 40°10' N. lat.	340.7	N/A	81.9	N/A	258.8
CANARY ROCKFISH a/ b/	Coastwide	106.8	N/A	56.9	N/A	49.9
COWCOD a/ c/	S of 40°10' N. lat.	4.0	N/A	1.4	N/A	2.6
DARKBLOTCHED ROCKFISH d/	Coastwide	317.2	95%	301.3	5%	15.9
PACIFIC OCEAN PERCH e/	N of 40°10' N. lat.	143.0	95%	135.9	5%	7.2
PETRALE SOLE a/	Coastwide	2,579.4	N/A	2,544.4	N/A	35.0
YELLOWEYE ROCKFISH a/	Coastwide	12.2	N/A	1.0	N/A	11.2
Arrowtooth flounder	Coastwide	3,410	95%	3,239	5%	170
Chilipepper	S of 40°10' N. lat.	1,604	75%	1,203	25%	401
Dover sole	Coastwide	48,406	95%	45,986	5%	2,420
English sole	Coastwide	9,640	95%	9,158	5%	482
Lingcod	N of 40°10' N. lat.	2,552	45%	1,148	55%	1,404
Lingcod	S of 40°10' N. lat.	995	45%	448	55%	547
Longnose skate a/	Coastwide	1,927	90%	1,734	10%	193
Longspine thornyhead	N of 34°27' N. lat.	3,124	95%	2,967	5%	156
Pacific cod	Coastwide	1,091	95%	1,036	5%	55
Pacific whiting	Coastwide	TBD	100%	TBD	0%	TBD
Sablefish	N of 36° N. lat.	0	See Table 1 c			
Sablefish	S of 36° N. lat.	1,714	42%	720	58%	994
Shortspine thornyhead	N of 34°27' N. lat.	1,686	95%	1,601	5%	84
Shortspine thornyhead	S of 34°27' N. lat.	881	NA	50	NA	831
Splitnose	S of 40°10' N. lat.	1,705	95%	1,619	5%	85
Starry flounder	Coastwide	1,524	50%	762	50%	762
Widow rockfish f/	Coastwide	1,880	91%	1,711	9%	169
Yellowtail rockfish	N of 40°10' N. lat.	5,560	88%	4,893	12%	667
Minor Shelf Rockfish complex a/	N of 40°10' N. lat.	1,872	60.2%	1,127	39.8%	745
Minor Shelf Rockfish complex a/	S of 40°10' N. lat.	1,575	12.2%	192	87.8%	1,383
Minor Slope Rockfish complex	N of 40°10' N. lat.	1,629	81%	1,319	19%	309
Minor Slope Rockfish complex	S of 40°10' N. lat.	673	63%	424	37%	249
Other Flatfish complex	Coastwide	8,545	90%	7,691	10%	855

a/ Allocations decided through the biennial specification process.

b/ 13.7 mt of the total trawl allocation of canary rockfish is allocated to the at-sea whiting fisheries, as follows: 5.7 mt for the mothership fishery, and 8.0 mt for the catcher/processor fishery.

c/ The cowcod fishery harvest guideline is further reduced to an ACT of 4.0 mt.

d/ Consistent with regulations at §660.55(c), 9 percent (27.1 mt) of the total trawl allocation for darkblotched rockfish is allocated to the whiting fisheries, as follows: 11.4 mt for the shorebased IFQ fishery, 6.5 mt for the mothership fishery, and 9.2 mt for the catcher/processor fishery. The tonnage calculated here for the whiting portion of the shorebased IFQ fishery contributes to the total shorebased trawl allocation, which is found at 660.140(d)(1)(ii)(D).

e/ Consistent with regulations at §660.55(c), 30 mt of the total trawl allocation for POP is allocated to the whiting fisheries, as follows: 12.6 mt for the shorebased IFQ fishery, 7.2 mt for the mothership fishery, and 10.2 mt for the catcher/processor fishery. The tonnage calculated here for the whiting portion of the shorebased IFQ fishery contributes to the total shorebased trawl allocation, which is found at 660.140(d)(1)(ii)(D).

f/ Consistent with regulations at §660.55(c), 500 mt of the total trawl allocation for widow rockfish is allocated to the whiting fisheries, as follows: 210 mt for the shorebased IFQ fishery, 120 mt for the mothership fishery, and 170 mt for the catcher/processor fishery. The tonnage calculated here for the whiting portion of the shorebased IFQ fishery contributes to the total shorebased trawl allocation, which is found at 660.140(d)(1)(ii)(D).