

Table 2a. To Part 660 Subpart C - 2012, and beyond, Specifications for OFL, ABC, ACL, ACT and Fishery Harvest guidelines (weights in metric tons).

Species	Area	OFL	ABC	ACL a/	ACT	Fishery HG
ROUND FISH:						
Lingcod	N of 42° N. lat. b/	2,251	2,151	2,151		1,880
	S of 42° N. lat. c/	2,597	2,164	2,164		2,157
Pacific Cod d/	Coastwide	3,200	2,222	1,600		1,200
Pacific Whiting e/	Coastwide	e/	e/	e/		135,481
Sablefish	N of 36° N. lat. f/	8,623	8,242	5,347	See Table 2c	
	S of 36° N. lat. g/			1,258		1,224
Cabezon	46°16' to 42° N. lat. h/	50	48	48		48
	S of 42° N. lat. i/	176	168	168		168
FLAT FISH:						
Dover sole j/	Coastwide	44,826	42,843	25,000		23,410
English sole k/	Coastwide	10,620	10,150	10,150		10,050
Petrale sole l/	Coastwide	1,279	1,222	1,160		1094.6
Arrowtooth flounder m/	Coastwide	14,460	12,049	12,049		9,971
Starry Flounder n/	Coastwide	1,813	1,511	1,360		1,353
Other flatfish o/	Coastwide	10,146	7,044	4,884		4,686
ROCK FISH:						
Pacific Ocean Perch p/	N of 40°10' N. lat.	1,007	962	183	157	144.1
Shorthelly q/	Coastwide	6,950	5,789	50		49
Widow r/	Coastwide	4,923	4,705	600		539.1
Canary s/	Coastwide	622	594	107		87
Chilipepper t/	S of 40°10' N. lat.	1,872	1,789	1,789		1,774
Bocaccio u/	S of 40°10' N. lat.	732	700	274		260.6
Splitnose v/	S of 40°10' N. lat.	1,610	1,538	1,538		1,531
Yellowtail w/	N of 40°10' N. lat.	4,573	4,371	4,371		3,872
Shortspine thornyhead x/	N of 34°27' N. lat.	2,358	2,254	1,556		1,511
	S of 34°27' N. lat.			401	359	
Longspine thornyhead y/	N of 34°27' N. lat.	3,483	2,902	2,064		2,020
	S of 34°27' N. lat.			366	363	
Cowcod z/	S of 40°10' N. lat.	13	10	3		2.7
Darkblotched aa/	Coastwide	497	475	296		277.3
Yelloweye bb/	Coastwide	48	46	17		11.1
California Scorpionfish cc/	S. of 34°27' N. lat.	132	126	126		124
Black	N of 46°16' N. lat. dd/	435	415	415		401
	S of 46°16' N. lat. ee/	1,169	1,117	1,000		1,000
Minor Rockfish North ff/ Nearshore Shelf Slope	Coastwide	3,820	3,414	2,227		2,116
	N of 40°10' N. lat.	116	99	99		99
		2,197	1,948	968		925
		1,507	1,367	1,160		1,092
Minor Rockfish South gg/ Nearshore Shelf Slope	Coastwide	4,291	3,712	2,341		2,290
	S of 40°10' N. lat.	1,145	990	990		990
		2,243	1,890	714		701
		903	832	626		599
SHARKS/SKATES/RATFISH/MORID						
Longnose Skate hh/	Coastwide	3,006	2,873	1,349		1,220
Other fish ii/	Coastwide	11,150	7,742	5,575		5,575

a/ ACLs and HGs are specified as total catch values. Fishery harvest guideline (HG) means the harvest guideline or quota after subtracting from the ACL of ACT any allocation for the Pacific Coast treaty Indian tribes, projected research catch, deductions for fishing mortality in non-groundfish fisheries, as necessary, and set-asides for EFPs.

b/ Lingcod north (Oregon and Washington). A new lingcod stock assessment was prepared in 2009. The lingcod north biomass was estimated to be at 62 percent of its unfished biomass in 2009. The OFL of 2,251 mt was calculated using an F_{MSY} proxy of $F_{45\%}$. The ABC of 2,151 mt was based on a 4 percent reduction from the OFL ($\sigma=0.36/P^*=0.45$) as it's a category 1 species. Because the stock is above $B_{40\%}$ coastwide, the ACL is set equal to the ABC. ACL is further reduced for the Tribal