

fishery (250 mt), resulting in a fishery HG of 1,880 mt.

c/ Lingcod south (California). A new lingcod stock assessment was prepared in 2009. The lingcod south biomass was estimated to be at 74 percent of its unfished biomass in 2009. The OFL of 2,597 mt was calculated using an F_{MSY} proxy of $F_{45\%}$. The ABC of 2,164 mt was based on a 17 percent reduction from the OFL ($\sigma=0.72/P^*=0.40$) as it's a category 2 species. Because the stock is above $B_{40\%}$ coastwide, the ACL is set equal to the ABC. An incidental open access set-aside of 7 mt is deducted from the ACL, resulting in a fishery HG of 2,157 mt.

d/ Pacific Cod. The 3,200 mt OFL is based on the maximum level of historic landings. The ABC of 2,222 mt is a 31 percent reduction from the OFL ($\sigma=1.44/P^*=0.40$) as it's a category 3 species. The 1,600 mt ACL is the OFL reduced by 50 percent as a precautionary adjustment. A set-aside of 400 mt is deducted from the ACL for the Tribal fishery, resulting in a fishery HG of 1,200 mt.

e/ Pacific whiting. The most recent stock assessment was prepared in January 2012. The 2012 Fishery Harvest Guideline (Fishery HG) is calculated as follows. U.S. TAC of 186,037 mt minus 48,556 mt for the Tribal allocation minus 2000 mt for catch in research activities and as non-groundfish bycatch, resulting in a fishery harvest guideline of 135,481 mt. The TAC for Pacific whiting is established under the provisions of the Pacific Hake/Whiting Agreement with Canada and the Pacific Whiting Act of 2006, 16 U.S.C. 7001-7010, and the international exception applies. Therefore, no OFL, ACL, or ACT values are provided for Pacific whiting.

f/ Sablefish north. A coastwide sablefish stock assessment was prepared in 2007. The coastwide sablefish biomass was estimated to be at 38.3 percent of its unfished biomass in 2007. The coastwide OFL of 8,623 mt was based on the 2007 stock assessment with a F_{MSY} proxy of $F_{45\%}$. The ABC of 8,242 mt is a 4 percent reduction from the OFL ($\sigma=0.36/P^*=0.45$) as it's a category 1 species. The 40-10 harvest policy was applied to the ABC to derive the coastwide ACL and then the ACL was apportioned north and south of 36° N. lat, using the average of annual swept area biomass (2003-2008) from the NMFS NWFSC trawl survey, between the northern and southern areas with 68 percent going to the area north of 36° N. lat. and 32 percent going to the area south of 36° N. lat. The northern portion of the ACL is 5,347 mt and is reduced by 535 mt for the tribal allocation (10 percent of the ACL north of 36° N. lat.) The 535 mt tribal allocation is reduced by 1.5 percent to account for discard mortality. Detailed sablefish allocations are shown in Table 2c.

g/ Sablefish South. That portion of the coastwide ACL (32 percent) apportioned to the area south of 36° N. lat. is 2,516 mt. An additional 50 percent reduction for uncertainty was made, resulting in an ACL of 1,258 mt. A set-aside of 34 mt is deducted from the ACL for EFP catch (26 mt), the incidental open access fishery (6 mt) and research catch (2 mt), resulting in a fishery HG of 1,224 mt.

h/ Cabezon (Oregon). A new cabezon stock assessment was prepared in 2009. The cabezon biomass in Oregon was estimated to be at 51 percent of its unfished biomass in 2009. The OFL of 50 mt was calculated using an F_{MSY} proxy of $F_{45\%}$. The ABC of 48 mt was based on a 4 percent reduction from the OFL ($\sigma=0.36/P^*=0.45$) as it's a category 1 species. Because the