

target year to rebuild of 2068 and an SPR rate of 82.7 percent. The amount anticipated to be taken during research activity is 0.1 mt and the amount expected to be taken during EFP activity is 0.2 mt, which results in a fishery HG of 2.7 mt.

aaa/ Darkblotched rockfish. A stock assessment update was prepared in 2009, based on the 2007 full assessment, and the stock was estimated to be at 27.5 percent of its unfished biomass in 2009. The OFL is projected to be 497 mt and is based on the 2009 stock assessment with an F_{MSY} proxy of $F_{50\%}$. The ABC of 475 mt is a 4 percent reduction from the OFL ($\sigma=0.36/P^*=0.45$) as it's a category 1 species. The ACL of 296 mt is based on a rebuilding plan with a target year to rebuild of 2025 and an SPR harvest rate of 64.9 percent. A set-aside of 18.7 mt is deducted from the ACL for the Tribal fishery (0.1 mt), the incidental open access fishery (15 mt), EFP catch (1.5) and research catch (2.1 mt), resulting in a fishery HG of 277.3 mt.

bbb/ Yelloweye rockfish. The stock was assessed in 2009 and was estimated to be at 20.3 percent of its unfished biomass in 2009. The 48 mt coastwide OFL was derived from the base model in the new stock assessment with an F_{MSY} proxy of $F_{50\%}$. The ABC of 46 mt is a 4 percent reduction from the OFL ($\sigma=0.36/P^*=0.45$) as it's a category 1 species. The 17 mt ACL is based on a rebuilding plan with a target year to rebuild of 2074 and an SPR harvest rate of 76 percent. A set-aside of 5.9 mt is deducted from the ACL for the Tribal fishery (2.3 mt), the incidental open access fishery (0.2 mt), EFP catch (0.1 mt) and research catch (3.3 mt) resulting in a fishery HG of 11.1 mt. Recreational HGs are being established as follows: Washington recreational, 2.6; Oregon recreational 2.4 mt; and California recreational 3.1 mt.

ccc/ California Scorpionfish south was assessed in 2005 and was estimated to be at 80 percent of its unfished biomass in 2005. The OFL of 132 mt is based on the new assessment with a harvest rate proxy of $F_{50\%}$. The ABC of 126 mt is a 4 percent reduction from the OFL ($\sigma=0.36/P^*=0.45$) as it's a category 1 species. Because the stock is above $B_{40\%}$, the ACL is set equal to the ABC. A set-aside of 2 mt is deducted from the ACL for the incidental open access fishery, resulting in a fishery HG of 124 mt.

ddd/ Black rockfish north (Washington). A stock assessment was prepared in 2007 for black rockfish north of 45°56' N. lat. (Cape Falcon, Oregon). The biomass in this area was estimated to be at 53 percent of its unfished biomass in 2007. The OFL from the assessed area is based on the 2007 assessment with a harvest rate proxy of $F_{50\%}$. The resulting OFL for the area north of 46°16' N. lat. (the Washington/Oregon border) is 435 mt, which is 97 percent of the OFL from the assessed area. The ABC of 415 mt for the area north of 46°16' N. lat. is a 4 percent reduction from the OFL ($\sigma=0.36/P^*=0.45$) as it's a category 1 species. The ACL was set equal to the ABC, since the stock is above $B_{40\%}$. A set-aside of 14 mt for the Tribal fishery results in a fishery HG of 401 mt.

eee/ Black rockfish south (Oregon and California). A 2007 stock assessment was prepared for black rockfish south of 45°56' N. lat. (Cape Falcon, Oregon) to the southern limit of the stock's distribution in Central California. The biomass in the south was estimated to be at 70 percent of its unfished biomass in 2007. The OFL from the assessed area is based on the 2007 assessment with a harvest rate proxy of $F_{50\%}$. Three