Table 2b. to Part 660, Subpart C – 2016, and Beyond, Allocations by Species or Species Group (weight in metric tons)

Species	Area	Fishery HG or ACT	Trawl		Non-trawl	
			%	Mt	%	Mt
BOCACCIO a/	S of 40°10' N. lat.	353.7	N/A	85.0	N/A	268.7
CANARY ROCKFISH a/ b/	Coastwide	109.8	N/A	58.5	N/A	51.3
COWCOD a/ c/	S of 40°10' N. lat.	4.0	N/A	1.4	N/A	2.6
DARKBLOTCHED ROCKFISH d/	Coastwide	325.2	95%	308.9	5%	16.3
PETRALE SOLE a/	Coastwide	2,673.4	N/A	2,638.4	N/A	35.0
PACIFIC OCEAN PERCH e/	N of 40°10' N. lat.	149.0	95%	141.6	5%	7.5
YELLOWEYE ROCKFISH a/	Coastwide	13.2	N/A	1.1	N/A	12.1
Arrowtooth flounder	Coastwide	3,241	95%	3,079	5%	162
Chilipepper	S of 40°10' N. lat.	1,595	75%	1,196	25%	399
Dover sole	Coastwide	48,406	95%	45,986	5%	2,420
English sole	Coastwide	6,991	95%	6,642	5%	350
Lingcod	N of 40'10° N. lat.	2,441	45%	1,098	55%	1,342
Lingcod	S of 40'10° N. lat.	937	45%	422	55%	515
Longnose skate a/	Coastwide	1,927	90%	1,734	10%	193
Longspine thornyhead	N of 34°27' N. lat.	2,969	95%	2,820	5%	148
Pacific cod	Coastwide	1,091	95%	1,036	5%	55
Pacific whiting	Coastwide	301,731	100%	301,731	0%	0
Sablefish	N of 36° N. lat.	0	See Table 1 c			
Sablefish	S of 36° N. lat.	1,875	42%	788	58%	1,088
Shortspine thornyhead	N of 34°27' N. lat.	1,667	95%	1,583	5%	83
Shortspine thornyhead	S of 34°27' N. lat.	871	NA	50	NA	821
Splitnose	S of 40°10' N. lat.	1,736	95%	1,649	5%	87
Starry flounder	Coastwide	1,529	50%	764	50%	764
Widow rockfish f/	Coastwide	1,880	91%	1,711	9%	169
Yellowtail rockfish	N of 40°10' N. lat.	5,314	88%	4,677	12%	638
Minor Shelf Rockfish complex a/	N of 40°10' N. lat.	1,880	60.2%	1,132	39.8%	748
Minor Shelf Rockfish complex a/	S of 40°10' N. lat.	1,576	12.2%	192	87.8%	1,384
Minor Slope Rockfish complex	N of 40°10' N. lat.	1,642	81%	1,330	19%	312
Minor Slope Rockfish complex	S of 40°10' N. lat.	675	63%	425	37%	250
Other Flatfish complex	Coastwide	7,039	90%	6,335	10%	704

a/ Allocations decided through the biennial specification process.

b/ 14.0 mt of the total trawl allocation of canary rockfish is allocated to the at-sea whiting fisheries, as follows: 5.8 mt for the mothership fishery, and 8.2 mt for the catcher/processor fishery.

c/ The cowcod fishery harvest guideline is further reduced to an ACT of 4.0 mt.

d/ Consistent with regulations at §660.55(c), 9 percent (27.8 mt) of the total trawl allocation for darkblotched rockfish is allocated to the whiting fisheries, as follows: 11.7 mt for the shorebased IFQ fishery, 6.7 mt for the mothership fishery, and 9.4 mt for the catcher/processor fishery. The tonnage calculated here for the whiting portion of the shorebased IFQ fishery contributes to the total shorebased trawl allocation, which is found at 660.140(d)(1)(ii)(D).

e/ Consistent with regulations at §660.55(c), 30 mt of the total trawl allocation for POP is allocated to the whiting fisheries, as follows: 12.6 mt for the shorebased IFQ fishery, 7.2 mt for the mothership fishery, and 10.2 mt for the catcher/processor fishery. The tonnage calculated here for the whiting portion of the shorebased IFQ fishery contributes to the total shorebased trawl allocation, which is found at 660.140(d)(1)(ii)(D).

f/ Consistent with regulations at §660.55(c), 500 mt of the total trawl allocation for widow rockfish is allocated to the whiting fisheries, as follows: 210 mt for the shorebased IFQ fishery, 120 mt for the mothership fishery, and 170 mt for the catcher/processor fishery. The tonnage calculated here for the whiting portion of the shorebased IFQ fishery contributes to the total shorebased trawl allocation, which is found at 660.140(d)(1)(ii)(D).