

Table 1b to Part 660, Subpart C –2017, Allocations by Species or Species Group (Weight in Metric Tons)

Species	Area	Fishery HG or ACT	Trawl		Non-trawl	
			Percent	Mt	Percent	Mt
BOCACCIO a/	S. of 40°10' N. lat.	774.6	39	302.4	61	472.2
COWCOD a/b/	S. of 40°10' N. lat.	4.0	36	1.4	64	2.6
DARKBLOTCHED ROCKFISH c/	Coastwide	563.8	95	535.6	5	28.2
PACIFIC OCEAN PERCH e/	N. of 40°10' N. lat.	231.6	95	220.0	5	11.6
YELLOWEYE ROCKFISH a/	Coastwide	14.6	NA	1.1	NA	13.1
Arrowtooth flounder	Coastwide	11,705.9	95	11,120.6	5	585.3
Big skate a/	Coastwide	436.6	95	414.8	5	21.8
Canary rockfish a/d/	Coastwide	1,466.6	NA	1,060.1	NA	406.5
Chilipepper	S. of 40°10' N. lat.	2,561.1	75	1,920.8	25	640.3
Dover sole	Coastwide	48,406.3	95	45,986.0	5	2,420.3
English sole	Coastwide	9,751.2	95	9,263.6	5	487.6
Lingcod	N. of 40°10' N. lat.	3,054.8	45	1,374.7	55	1,680.2
Lingcod	S. of 40°10' N. lat.	1,242.0	45	558.9	55	683.1
Longnose skate a/	Coastwide	1,853.0	90	1,667.7	10	185.3
Longspine thornyhead	N. of 34°27' N. lat.	2,847.2	95	2,704.8	5	142.4
Pacific cod	Coastwide	1,091.0	95	1,036.4	5	54.5
Pacific whiting f/	Coastwide	362,682.0	100	362,682.0	0	0.0
Petrale sole	Coastwide	2,895.1	95	2,750.3	5	144.8
Sablefish	N. of 36° N. lat.	N/A	See Table 1c			
Sablefish	S. of 36° N. lat.	1,859.0	42	780.8	58	1,078.2
Shortspine thornyhead	N. of 34°27' N. lat.	1,654.0	95	1,571.3	5	82.7
Shortspine thornyhead	S. of 34°27' N. lat.	863.7	NA	50.0	NA	813.7
Splitnose rockfish	S. of 40°10' N. lat.	1,749.3	95	1,661.8	5	87.5
Stary flounder	Coastwide	1,271.7	50	635.9	50	635.9
Widow rockfish g/	Coastwide	13,290.3	91	12,094.2	9	1,196.1
Yellowtail rockfish	N. of 40°10' N. lat.	5,166.1	88	4,546.1	12	619.9
Minor Shelf Rockfish a/	N. of 40°10' N. lat.	1,965.2	60	1,183.1	40	782.1
Minor Slope Rockfish	N. of 40°10' N. lat.	1,689.9	81	1,368.8	19	321.1
Minor Shelf Rockfish a/	S. of 40°10' N. lat.	1,575.8	12	192.2	88	1,383.6
Minor Slope Rockfish	S. of 40°10' N. lat.	686.8	63	432.7	37	254.1
Other Flatfish	Coastwide	8,306.0	90	7,475.4	10	830.6

a/ Allocations decided through the biennial specification process.

b/ The cowcod fishery harvest guideline is further reduced to an ACT of 4.0 mt.

c/ Consistent with regulations at §660.55(c), 9 percent (48.2 mt) of the total trawl allocation for darkblotched rockfish is allocated to the Pacific whiting fishery, as follows: 20.2 mt for the Shorebased IFQ Program, 11.6 mt for the MS sector, and 16.4 mt for the C/P sector. In July 2017, the amounts available to the mothership and catcher/processor fisheries were raised by 25 mt, to 36.6 mt for the mothership fishery and to 41.4 mt for the catcher/processor fishery, by distributing equally the full 50 mt initially deducted from the ACL to account for unforeseen catch events, consistent with §660.60(c)(3)(ii). The tonnage calculated here for the Pacific whiting IFQ fishery contributes to the total shorebased trawl allocation, which is found at §660.140(d)(1)(ii)(D).

d/ Canary rockfish is allocated approximately 72 percent to trawl and 28 percent to non-trawl. 46 mt of the total trawl allocation of canary rockfish is allocated to the MS and C/P sectors, as follows: 30 mt for the MS sector, and 16 mt for the C/P sector.

e/ Consistent with regulations at §660.55(c), 17 percent (37.4 mt) of the total trawl allocation for POP is allocated to the Pacific whiting fishery, as follows: 15.7 mt for the Shorebased IFQ Program, 9.0 mt for the MS sector, and 12.7 mt for the C/P sector. In May 2017, the amounts available to the mothership and catcher/processor fisheries were raised by 3.5 mt, to 12.5 mt for the mothership fishery and to 16.2 mt for the catcher/processor fishery, by distributing 7.0 mt of the 10 mt initially deducted from the ACL to account for mortality in the incidental open access fishery, consistent with §660.60(c)(3)(ii). In July 2014, the amounts available to the mothership and catcher processor fisheries were each raised by 12.5 mt, to 25 mt for the mothership fishery and to 28.7 mt for the catcher/processor fishery, by distributing equally the full 25 mt initially deducted from the ACL to account for unforeseen catch events, consistent with §660.60(c)(3)(ii). The tonnage calculated here for the Pacific whiting IFQ fishery contributes to the total shorebased trawl allocation, which is found at §660.140(d)(1)(ii)(D).

f/ Consistent with regulations at §660.55(f), the commercial harvest guideline for Pacific whiting is allocated as follows: 34 percent (123,312 mt) for the C/P Coop Program; 24 percent (87,044 mt) for the MS Coop Program; and 42 percent (152,326.5 mt) for the Shorebased IFQ Program. No more than 5 percent of the Shore based IFQ Program allocation (7,616 mt) may be taken and retained south of 42° N. lat. before the start of the primary Pacific whiting season north of 42° N. lat.

g/ Consistent with regulations at §660.55(c), 10 percent (1,209.4 mt) of the total trawl allocation for widow rockfish is allocated to the whiting fisheries, as follows: 508.0 mt for the shorebased IFQ fishery, 290.3 mt for the mothership fishery, and 411.2 mt for the catcher/processor fishery. The tonnage calculated here for the whiting portion of the shorebased IFQ fishery contributes to the total shorebased trawl allocation, which is found at §660.140(d)(1)(ii)(D).