

kk/ Cabezon/kelp greenling (Washington). 2 mt is deducted from the ACL to

accommodate the Tribal fishery, therefore the fishery HG is 18 mt.

ll/Nearshore Rockfish north of 40°10' N lat. 3.08 mt is deducted from the ACL to accommodate the Tribal fishery (1.5 mt), EFP catch (0.5 mt), research (0.47 mt), and the incidental open access fishery (0.61 mt), resulting in a fishery HG of 75.9 mt. State specific HGs are Washington (18.4 mt), Oregon (22.7 mt), and California (37.6 mt).

mm/ Nearshore Rockfish south of 40°10' N lat. 4.42 mt is deducted from the ACL to accommodate research catch (2.68 mt) and the incidental open access fishery (2.68 mt), resulting in a fishery HG of 1,011.6 mt.

nn/ Other Fish. The Other Fish complex is comprised of kelp greenling off California and leopard shark coastwide. 21.34 mt is deducted from the ACL to accommodate EFP catch (0.1 mt), research (6.29 mt), and the incidental open access fishery (14.95 mt), resulting in a fishery HG of 201.7 mt.

oo/ Other Flatfish. The Other Flatfish complex is comprised of flatfish species managed in the PCGFMP that are not managed with stock-specific OFLs/ABCs/ACLs. Most of the species in the Other Flatfish complex are unassessed and include: butter sole, curlfin sole, flathead sole, Pacific sanddab, rock sole, sand sole, and rex sole. 220.89 mt is deducted from the ACL to accommodate the Tribal fishery (60 mt), EFP catch (0.1 mt), research (23.63 mt), and the incidental open access fishery (137.16 mt), resulting in a fishery HG of 4,581.1 mt.

pp/ Shelf Rockfish north of 40°10' N lat. 72.44 mt is deducted from the ACL to accommodate the Tribal fishery (30 mt), EFP catch (1.5 mt), research (15.32 mt), and the incidental open access fishery (25.62 mt), resulting in a fishery HG of 1,438.66 mt.

qq/ Shelf Rockfish south of 40°10' N lat. 132.77 mt is deducted from the ACL to accommodate EFP catch (50 mt), research catch (15.1 mt), and the incidental open access fishery (67.67 mt) resulting in a fishery HG of 1,305.2 mt.