

ACL to accommodate the Tribal fishery (60 mt), EFP catch (0.1 mt), research (23.63 mt), and

the incidental open access fishery (137.16 mt), resulting in a fishery HG of 4,617.1 mt.

pp/ Shelf Rockfish north of 40°10' N lat. 72.44 mt is deducted from the ACL to

accommodate the Tribal fishery (30 mt), EFP catch (1.5 mt), research (15.32 mt), and the

incidental open access fishery (25.62 mt), resulting in a fishery HG of 1,377.6 mt.

qq/ Shelf Rockfish south of 40°10' N lat. 132.77 mt is deducted from the ACL to

accommodate EFP catch (50 mt), research catch (15.1 mt), and the incidental open access fishery

(67.67 mt) resulting in a fishery HG of 1,295.2 mt.

rr/ Slope Rockfish north of 40°10' N lat. 65.89 mt is deducted from the ACL to

accommodate the Tribal fishery (36 mt), EFP catch (1.5 mt), and research (10.51 mt), and the

incidental open access fishery (18.88 mt), resulting in a fishery HG of 1,502.1 mt.

ss/ Slope Rockfish south of 40°10' N lat. 38.94 mt is deducted from the ACL to

accommodate EFP catch (1 mt), and research (18.21 mt), and the incidental open access fishery

(19.73 mt), resulting in a fishery HG of 666.1 mt. Blackgill rockfish has a stock-specific HG for

the entire groundfish fishery south of 40°10' N lat. set equal to the species' contribution to the 40-

10-adjusted ACL. Harvest of blackgill rockfish in all groundfish fisheries south of 40°10' N lat.

counts against this HG of 174 mt.