

**Table 2b. to Part 660, Subpart C—2022, and Beyond, Allocations by Species or Species Group [Weight in Metric Tons]**

Stocks/Stock Complexes	Area	Fishery HG or ACT a/ b/	Trawl		Non-Trawl	
			%	Mt	%	Mt
YELLOWEYE ROCKFISH <sup>a/</sup>	Coastwide	42.2	8	3.4	92	38.8
Arrowtooth flounder	Coastwide	6,362.9	95	6,044.8	5	318.1
Big skate <sup>a/</sup>	Coastwide	1,331.7	95	1,265.1	5	66.6
Bocaccio <sup>a/</sup>	S of 40°10' N. lat.	1,676.2	39.04	654.4	60.96	1,021.8
Canary rockfish <sup>a/</sup>	Coastwide	1,237.6	72.281	894.6	27.719	343.1
Chilipepper rockfish	S of 40°10' N. lat.	2,161.3	75	1,621	25	540.3
Cowcod <sup>a/</sup>	S of 40°10' N. lat.	50	36	18	64	32
Darkblotched rockfish	Coastwide	811.9	95	771.3	5	40.6
Dover sole	Coastwide	4,8402.8	95	45,982.7	5	2,420.1
English sole	Coastwide	8,850.4	95	8,407.8	5	442.5
Lingcod	N of 40°10° N. lat.	4,679.6	45	2,105.8	55	2,573.8
Lingcod <sup>a/</sup>	S of 40°10° N. lat.	1,159	40	463.6	60	695.4
Longnose skate <sup>a/</sup>	Coastwide	1,509.6	90	1,358.6	10	151
Longspine thornyhead	N of 34°27' N. lat.	2,398.3	95	2,278.4	5	119.9
Pacific cod	Coastwide	1,093.9	95	1,039.2	5	54.7
Pacific ocean perch	N of 40°10' N. lat.	3,686.3	95	3,502	5	184.3
Pacific whiting <sup>c/</sup>	Coastwide	TBD	100	TBD	0	0
Petrale sole <sup>a/</sup>	Coastwide	3,272.5	-	3,242.5	-	30
Sablefish	N of 36° N. lat.	NA	See Table 1c			
Sablefish	S of 36° N. lat.	1,781.6	42	748.3	58	1,033.3
Shortspine thornyhead	N of 34°27' N. lat.	1,314.6	95	1,248.9	5	65.7
Shortspine thornyhead	S of 34°27' N. lat.	730.3		50		680.3
Splitnose rockfish	S of 40°10' N. lat.	1,611.6	95	1,531	5	80.6
Starry flounder	Coastwide	343.6	50	171.8	50	171.8
Widow rockfish <sup>a/</sup>	Coastwide	13,539.7		13,139.7		400
Yellowtail rockfish	N of 40°10' N. lat.	4,793.5	88	4,218.2	12	575.2
Other Flatfish	Coastwide	4,617.1	90	4,155.4	10	461.7
Shelf Rockfish <sup>a/</sup>	N of 40° 10' N. lat.	1,377.6	60.2	829.3	39.8	548.3
Shelf Rockfish <sup>a/</sup>	S of 40° 10' N. lat.	1,295.2	12.2	158	87.8	1,137.2
Slope Rockfish	N of 40° 10' N. lat.	1,502.1	81	1,216.7	19	285.4
Slope Rockfish <sup>a/</sup>	S of 40° 10' N. lat.	666.1		523.9		142.2

a/ Allocations decided through the biennial specification process.

b/ The cowcod fishery harvest guideline is further reduced to an ACT of 50 mt.

c/ Consistent with regulations at §660.55(i)(2), the commercial harvest guideline for Pacific whiting is allocated as follows: 34 percent for the C/P Coop Program; 24 percent for the MS