Species/Species Group/Area	Trawl Allocation (mt)	At-sea Whiting set asides	Shorebased IFQ			
			Non-Whiting		Whiting	
			percent	mt	percent	mt
Pacific Ocean Perch	137	17.4 (10.2 catcher/processor + 7.2 mothership)	89.5% [Remaining]	107	10.5% (Greater of 17% or 30 mt to shorebased + at-sea whiting)	12.6
Widow rockfish	491	147.9 (86.7 catcher/processor + 61.2 mothership)	68.7% (Remaining)	235	31.3% (52% to shorebased + at-sea whiting)	107.1
Yellowtail rockfish North of 40°10' N. lat.	3,401	300	90.3% (Remaining)	2,801	9.7% (300 mt)	300
Darkblotched rockfish	265	14.5 (8.5 catcher/ processors + 6 mothership)	95.8% (Remaining)	240	4.20% (Greater of 9% or 25 mt to shorebased + at-sea whiting)	10.5
Minor slope rockfish	377	na	100%	377	0.0%	0
South of 40°10' N. lat.						
Minor shelf rockfish						
North of 40°10' N. lat.	557	35	82.6%	431.2	17.4%	90.8
South of 40°10' N. lat.	86	na	100%	86	0.0%	0
Canary Rockfish	34.1	8.2 (4.8 catcher/processor + 3.4 mothership)	77.2%	20	22.8%	5.9
Bocaccio	60	na	100%	60	0.0%	0
Cowcod	1.8	na	100%	1.8	0.0%	0
Yelloweye Rockfish	0.6	0	100%	0.6	0.0%	0