

Title:
9 pt. Helvetica Bold
Italic, left justified

Body text:
6 pt. Helvetica Regular with
6.5 pts. leading, left justified

Bullet: 5 pt.
Solid square

Subheadings:
6 pt. Helvetica Bold,
left justified

Headings:
8 pt. Helvetica Bold
Italic, left justified

Drug Facts

Active ingredients (in each tablet) Purpose

Aluminum hydroxide gel 200 mg.....	Antacid
Magnesium hydroxide 200 mg.....	Antacid
Simethicone 25 mg.....	Antigas

Uses

- relieves symptoms referred to as gas
- relieves: ■ heartburn ■ acid indigestion ■ sour stomach
 ■ upset stomach due to these symptoms

Warnings

Ask a doctor before use if you have kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last for more than 2 weeks

Keep out of reach of children.

Directions ■ chew 1 to 4 tablets 4 times daily

- do not take more than 16 tablets in 24 hours or use the maximum dosage for more than 2 weeks

Inactive ingredients D&C red no. 30, D&C yellow no. 10, dextrose, FD&C blue no. 1, glycerin, magnesium stearate, mannitol, saccharin sodium, sorbitol, starch, sugar, talc

Right justified

2.5 point barline

0.5 point hairline

Bulleted information may start on same line as headings (except Warnings) and subheadings and need not be vertically aligned

Dark type on light background

Box barline omitted; color contrast used to highlight Drug Facts information